How (Not) to Integrate Blood Subtyping Technology to Kidney Exchange

Tayfun Sönmez M. Utku Ünver Özgür Y lmaz

October 2017

Abstract

Even though kidney exchange became an important source of kidney transplants over the

preliminaries, In Section 3 we introduce the model, and in Section 4 we analyze the impact of availability and timing of various blood subtyping technologies on the number of living-donor kidney transplants. In Section 5, we show via computational simulations that all our theoretical results are veri ed using kidney-donation data from the US. In Section 6, we conclude. The Appendix is devoted to proofs (Appendix A), the extended analysis allowing for 3-way kidney exchange in addition to 2-way exchange (Appendix B), additional results related to simulations (Appendix C).

it is possible to calculate the PRA distribution using di erent data sets for various types of individuals; for example, waitlisted kidney patients using UNOS data).

2.2 Blood-Type Compatibility

Blood-type compatibility is the more structured of the two primary medical requirements for kidney transplantation. The standard blood type classi cation is known as ABO grouping. Under this classi cation there are two types of red blood cell proteins (or antigens) referred to as antigen A and antigen B. An individual can have any combination of these two antigens, and individuals produce antibodies against the antigens they lack. There are four blood types determined by the presence or absence of these two antigens on the surface of red blood cells:

1. Blood type O: Has neither A nor B antigens on red cells (but both antibody anti-A and antibody anti-B are in the plasma).
2. Blood type A: Has only the A antigen on red cells (and antibody anti-B in the plasma).
3. Blood type B: Has only the B antigen on red cells (and antibody anti-A in the plasma).
4. Blood type AB: Has both A and B antigens on red cells (but neither antibody anti-A nor antibody anti-B in the plasma).

The above-described blood type classi cation induces the standard ABO blood-type compatibility , ory bloo

2.3 Blood Subtype A2/A2B Compatibility

Unlike the red blood cell antigen B, antigen A is not a single antigen and it consists of two major subtypes, A1 and A2. When transplanted to a blood type B or O patient, a subtype A2 kidney generates considerably weaker antibody response than a subtype A1 kidney. Similarly, subtype A2B kidneys (those with both subtype A2 and type B antigens) generate weaker antibody response than A1B kidneys (those with both subtype A1 and type B antigens) in blood type B patients. The antibody response to subtype A2 (or A2B) kidneys is not uniform across all

under the baseline scenario. Patients who are both tissue-type and blood-type compatible with their donors receive a kidney transplant directly from their compatible donors. In this baseline scenario, patients who are either tissue-type incompatible or blood-type incompatible are transferred to a single kidney-exchange pool. This process determines the composition of the kidney-exchange pool.

In addition to the baseline ABO-compatibility technology, we also consider the three subtyping technologies A2-to-B compatibility, A2-to-O compatibility, and full compatibility, described in Section 2.3. When available, a subtyping technology will replace the baseline ABO-compatibility technology, thereby potentially removing a barrier to direct donation. Therefore compared to the baseline ABO compatibility, the availability of each of the three subtyping technologies will potentially extend the set of patients who receive a transplant via direct donation and potentially shrink the set of patients who are transferred to the kidney-exchange pool.

A 2-way kidney exchange involves two patients, each of whom is compatible with the other patient's donor. When a 2-way exchange is carried out, both patients receive a kidney from the other patient's donor. Once the kidney-exchange pool forms under any of the above-described bloodtyping/subtyping technologies, a maximal-size kidney exchange is determined for the given pool of incompatible patient donor pairs as in Roth, Sönmez, and Ünver (2005b, 2007). Our focus is to analyze the impact of availability of various subtyping technologies on the total number of living-donor transplants, including direct transplants from patients' compatible donors and transplants from kidney exchanges.

Following Roth, Sönmez, and Ünver (2007), we rely on the following three assumptions about the composition of the kidney-exchange pool to derive our analytical results. While these assumptions are used for our analytical results, they will be dispensed with in our simulations in Section 5. We will observe that all qualitative implications of our analytical results are veri ed by our simulations in the absence of these assumptions. We next motivate and formally state these assumptions.

Let X, Y be two distinct blood types such that a blood type Y donor can donate to a blood type X patient. Being blood-type compatible, a pair of type X-Y is only transferred to the kidney-exchange pool when the pair is tissue-type incompatible, a relatively rare event. In contrast, a pair of the opposite type Y-X is always transferred to the kidney-exchange pool because they are blood-type incompatible. Therefore, in the long run, there will be an abundance of Y-X pairs in the kidney-exchange pool in comparison with X-Y pairs, and thus it will not be possible to match all pairs of type Y-X.[9] We refer to this situation as a pair of

[9]That is why there is an abundance of blood type O patients with non-O donors in kidney-exchange pools all over the world.

5.2 Simulation Results

We run S = 500 2-way-exchange simulations and report the results as averages and sample standard deviations in Table 2.[17]

Under A2-to-B compatibility when subtyping tests are conducted before joining the pool, the average number of transplants achievable through 2-way exchange (and direct donation if needed) decreases from Scenario 1 the exchange practice under ABO compatibility only to Scenario 2(a)i the de facto exchange policy with A2 compatibility as predicted by Proposition 1. This decrease is statistically signi cant with a paired t-test for degrees of freedom 499 with t-statistic 4.3366 that leads to a p-value less than 10^{-5}.

We also document how the e ect of this transition is shared across patients of di erent ethnic backgrounds (see Table 3). We observe that for whites, there is a decrease, which is statistically signi cant (with a paired t-test p-value of less than 10^{-5}). For other races, changes are not statistically signi cant.

The direction of other changes across all treatments are in accordance with the theoretical results of Section 4, as seen in Table 2. In particular, the average number of transplants signi cantly increases from Scenario 1 and 2(a)i to 2(a)ii and 2(a)iii (cf. Proposition 2), while there is a signi cant decrease from Scenario 2(a)ii to 2(a)iii (cf. Proposition 3). The same pattern is true for all races (see Table 3). This corresponds to an 11.05 13.74 percent increase from A2-to-B compatibility to A2-to-O compatibility (from Scenario 2(a)i to 2(a)ii) for di erent population sizes. Moreover, when the A2 subtype transplantation is done uncoordinated with exchange before the exchange participation decision, even full compatibility brings a substantial decrease to the number of transplants: a 2 percent decrease is observed in the number of transplants with respect to A2-to-O compatibility (from Scenario 2(a)ii to 2(a)iii).

When the subtyping tests are conducted after joining the exchange pool, the corresponding numbers of transplants increase with respect to the cases when subtyping tests are conducted before joining the pool (the signs and statistical signi cance of di erences across scenarios are as predicted by the theoretical results). What is striking is that the maximum average number of transplants under optimal 2-way exchange and live donation (Scenario 2(b)iii), i.e., the maximum achievable, is not too much di erent (although the increase is highly statistically signi cant) from the A2-to-O compatibility scenario with subtyping before joining the pool (Scenario 2(a)ii): the di erence is only between 0.7 2.5 percent across di erent population sizes. If A2-to-B compatibility could totally be eliminated for live donation and we could only use A2-to-O compatibility, even if we could not defer subtype testing until pairs join the exchange pool, we would lose very little with respect to the optimal policy recommendation.

[17] In Appendix C, we report the results of simulations for 2&3-way technology in Tables 5 and 7.

Simulation averages and sample standard deviations of S = 500 simulations			
	2-way exchanges		
Incomp.	1. Under ABO	2. Under A2 compatibility	

0.5580

Races	Number of pairs		Number of patients of each race matched under 2-way exchanges							
			1. Under	2. Under A2 compatibility						
				A subtype test timing: (a) Before joining exchange pool			A subtype test timing: (b) After joining exchange pool			
	Comp. (under ABO compatibility)	Inc.	ABO comp.	i. A2/A2B ! B only	ii. A2 ! O only	iii. A2/A2B ! B and A2 ! O	i. A2/A2B ! B only	ii. A2 ! O only	iii. A2/A2B ! B and A2 ! O
White	41.4280 (5.2122)	40.0520 (5.0071)	13.0340 (4.2869)	13.3360 (4.4070)	14.8540 (4.7478)	14.8480 (4.7360)	13.5760 (4.4569)	14.9380 (4.7589)	15.2460 (4.7958)
Black	6.3380 (2.4074)	6.4560 (2.4609)	2.0960 (1.4857)	2.1840 (1.5527)	2.4100 (1.6270)	2.3960 (1.6311)	2.2060 (1.5609)	2.4100 (1.6258)	2.4720 (1.6718)
Asian	2.4540 (1.4860)	2.6800 (1.6562)	0.9940 (1.0641)	1.0260 (1.0750)	1.0960 (1.1159)	1.0900 (1.0974)	1.0640 (1.1001)	1.0900 (1.1137)	1.1180 (1.1414)
American Indian	0.1880 (0.4207)	0.2120 (0.4329)	0.0680 (0.2520)	0.0700 (0.2554)	0.0840 (0.2848)	0.0820 (0.2746)	0.0700 (0.2554)	0.0880 (0.2836)	0.0800 (0.2716)
Paci c Islander	0.1000 (0.3003)	0.0920 (0.2962)	0.0240 (0.1532)	0.0220 (0.1468)	0.0320 (0.1762)	0.0300 (0.1708)	0.0240 (0.1532)	0.0280 (0.1651)	0.0280 (0.1651)
TOTAL out of n = 100 pairs	50.508 (5.1942)	49.492 (5.1942)	16.216 (4.9236)	16.638 (5.0666)	18.476 (5.4517)	18.446 (5.4113)	16.940 (5.1265)	18.554 (5.450)	18.944 (5.5352)
White	207.8560 (11.8609)	198.8320 (11.3236)	72.1760 (9.6702)	72.7000 (10.0259)	82.6220 (10.9621)	81.3440 (10.7693)	74.4220 (10.0918)	83.0460 (10.8819)	83.9060 (10.9744)
Black	31.5340 (5.4129)	32.7300 (5.4398)	12.4280 (3.6856)	12.7680 (3.6771)	14.2040 (3.9595)	13.9880 (3.8618)	12.9220 (3.7449)	14.1240 (3.9288)	14.3940 (3.9035)
Asian	12.5540 (3.3295)	13.2520 (3.6572)	5.4940 (2.4236)	5.6740 (2.4641)	6.1420 (2.5516)	6.1480 (2.5907)	5.8260 (2.5394)	6.1500 (2.5052)	6.2560 (2.5713)
American Indian	1.0680 (1.0685)	1.0200 (1.0206)	0.3300 (0.6244)	0.3240 (0.6097)	0.3820 (0.6730)	0.3480 (0.6290)	0.3340 (0.6255)	0.3740 (0.6442)	0.3740 (0.6220)
Paci c Islander	0.5580 (0.7535)	0.5960 (0.7307)	0.2080 (0.4485)	0.2160 (0.4624)	0.2500 (0.4940)	0.2420 (0.4857)	0.2160 (0.4536)	0.2580 (0.5099)	0.2300 (0.4749)
TOTAL	253.570	246.430							

Analytically predicted averages and standard deviations of S = 500 simulated pools					
	2-way exchanges				
	1. Under	2. Under A2 compatibility			
Incomp.	ABO	A subtype	A2 compatibility technology		
pairs	compatibility	test	i. A2/A2B ! B	ii. A2 ! O	iii. A2/A2B ! B
		out of	only	only	and A2 ! O
49.420		(a) Before joining	15.906	19.640	19.238
(5.1942)	16.308	exchange pool	(4.8514)	(5.3778)	(5.249)
out of	(4.9778)	(b) After			

sizes. Thus, A2/A2B ! B technology results with slightly less reduction in number of transplants than predicted by analytical results.

These two e ects work in opposite directions when both A2/A2B B and A2 ! O technologies are available. The second e ect is smaller than the rst e ect. Thus, most of our analytical predictions in terms of the directions of the changes remain intact through the simulations.

2&3-way simulations are very similar, as reported in Tables 5, 7, and 6, respectively, in Appendix C. There is a slight di erence in the case of 2&3-way exchanges: As shown in Appendix B (and the intuition explained in Section 4.3), the e ciency loss due to B-A2 pairs not participating in the exchange is compensated for certain distributions of the pairs in the exchange pool. As a result, the analytically predicted average number of transplants does not decrease when A2-to-B compatibility is integrated (Table 6), and the gap between simulations and analytical results is even smaller compared to only 2-way exchanges (Tables 5 and 6).

6 Conclusion

Roth, Alvin E., Tayfun Sönmez, and M. Utku Ünver (2005c), "Transplant center incentives in kidney exchange." Harvard University and Koç University, unpublished mimeo.

Roth, Alvin E., Tayfun Sönmez, and M. Utku Ünver (2007), "Efficient kidney exchange: Coincidence of wants in markets with compatibility-based preferences." *American Economic Review*, 97, 828–851.

Sönmez, Tayfun (2013), "Bidding for army career specialties: Improving the ROTC branching mechanism." *Journal of Political Economy*, 121, 186–219.

Sönmez, Tayfun and M. Utku Ünver (2014), "Altruistically unbalanced kidney exchange." *Journal of Economic Theory*, 152, 105–129.

Sönmez, Tayfun and M. Utku Ünver (2015), "Enhancing the efficiency of and equity in transplant organ allocation via incentivized exchange." Available as Boston College Working Paper 868.

Sönmez, Tayfun, M. Utku Ünver, and M. Bumin Yenmez (2017), "Incentivized kidney exchange." Working paper.

Terasaki, Paul I., David W. Gjertson, and J. Michael Cecka (1998), "Paired kidney exchange is not a solution to ABO incompatibility." *Transplantation*, 65.

Toulis, Panos and David C. Parkes (2015), "Design and analysis of multi-hospital kidney exchange mechanisms using random graphs." *Games and Economic Behavior*, 91, 360–382.

Ünver, M. Utku (2010), "Dynamic kidney exchange." *Review of Economic Studies*, 77, 372–414.

Zenios, Stefanos A., E. Steve Woodle, and Lainie Friedman Ross (2001), "Primum non nocere: Avoiding increased waiting times for individual racial and blood-type subsets of kidney wait list candidates in a living donor/cadaveric donor exchange program." *Transplantation*, 72, 648–654.