



# Exposure to Bisphenol A and diabetes risk in Mexican women

Luke Murphy<sup>1,2</sup>; Ángel Mérida-Ortega<sup>1</sup>; Mariano E. Cebrán<sup>3</sup>; Lucia Hernández-Garcia diego<sup>4</sup>; Humberto Gómez-Ruiz<sup>4</sup>; Brenda Gamboa-Loira<sup>1</sup>; Lizbeth López-Carrillo<sup>1</sup>

<sup>1</sup>Center for Research in Population Health, National Institute of Public Health, Cuernavaca, Morelos, México <sup>2</sup>Department of Biology, Boston College, Chestnut Hill, MA, US <sup>3</sup>Department of Toxicology, CINVESTAV, Ciudad de México, México <sup>4</sup>Department of Analytical Chemistry, National Autonomous University of Mexico, Ciudad de México, México



BPA is an endocrine disrupting chemical (EDC) used in the production of polycarbonate plastics and epoxy resins. As one of the most widely produced chemicals in the world, BPA has been linked to various health problems, including diabetes. In this study, we investigated the relationship between BPA exposure and diabetes risk in Mexican women. We found that women with higher BPA levels were more likely to have diabetes compared to those with lower levels. This suggests that BPA may contribute to the development of diabetes in women.

